

REVIEW ARTICLE

Soluble syndecans: biomarkers for diseases and therapeutic options

Correspondence Jessica Bertrand, Department of Orthopaedic Surgery, Otto von Guericke University Magdeburg, Leipziger Straße 44, Magdeburg 39120, Germany. E-mail: jessica.bertrand@med.ovgu.de

Received 13 March 2018; **Revised** 1 June 2018; **Accepted** 8 June 2018

Jessica Bertrand  and Miriam Bollmann

Department of Orthopaedic Surgery, Otto von Guericke University Magdeburg, Magdeburg, Germany

Syndecans are important mediators of signalling by transmitting external stimuli into the cells. This role in signal transduction has been attributed mainly to the membrane-bound syndecans. In the last years, however, the soluble ectodomain of syndecans generated by shedding has come into the focus of research as this process has been shown to modulate the syndecan-dependent signalling pathways, as well as other pathways. This review summarizes the current knowledge about the induction of syndecan shedding and the different pathways modulated by shed syndecan proteins. This review summarizes the known and putative sheddases for each syndecan and describes the exemplary conditions of sheddase activity for some syndecans. This review summarizes the proposed use of shed syndecans as biomarkers for various diseases, as the shedding process of syndecans depends crucially on tissue- and disease-specific activation of the sheddases. Furthermore, the potential use of soluble syndecans as a therapeutic option is discussed, on the basis of the current literature.

LINKED ARTICLES

This article is part of a themed section on Translating the Matrix. To view the other articles in this section visit <http://onlinelibrary.wiley.com/doi/10.1111/bph.v176.1/issuetoc>

Abbreviations

ADAMs, a disintegrin and metalloproteinase; ECM, extracellular matrix; FAK, focal adhesion kinase; GAG, glycosaminoglycan; HS, heparan sulfate; Sdc, syndecan; TACE, TNF- α -converting enzyme, ADAM-17; VEGFR2, VEGF receptor 2

Syndecan structure

Syndecans belong to the family of type I transmembrane heparan sulfate proteoglycans, which consist of four members in vertebrates (**Sdc1**, **Sdc2**, **Sdc3** and **Sdc4**). The core protein of these proteoglycans is composed of an extracellular, transmembrane and intracellular domain (Figure 1). Sdc1, Sdc2 and Sdc4 are translated with a signal peptide, which is cleaved during the processing of the protein (Figure 1). All syndecans span the plasma membrane via a 24–25 amino acid long hydrophobic transmembrane domain. The transmembrane domain includes a GXXXG motif, which allows for a strong, SDS-resistant homodimerization of syndecans (Choi *et al.*, 2005). The transmembrane and cytoplasmic domains share approximately 60–70% amino acid sequence identity between all family members (David, 1993).

The membrane proximal region C1 is highly conserved among all syndecans (90%) and also among different species, as well as the C-terminal C2 region (100% conservation between syndecans). These two domains flank a variable region V1, which differs for each syndecan and exhibits only 15% similarity between all syndecans (Figure 1). However, Sdc1 and Sdc3 have a higher similarity in this variable region, as well as Sdc2 and Sdc4. This finding gave rise to the assumption that these members of the syndecan family are more closely related to each other than to the others (Carey, 1997).

The ectodomains of syndecans share only weak homology between all four members (Figure 1). The putative glycosaminoglycan (GAG) attachment sites have similar consensus sequences. Two attachment sites have the consensus sequence SGXG, and three others have the consensus sequence (E/D) GSG (E/D). The existence of GAG binding sites either at both ends of the ectodomain (Sdc1 and Sdc3) or only at the distal part (Sdc2 and Sdc4) of the ectodomain is another indicator for the sub-classification of syndecans (Elenius and Jalkanen, 1994). Syndecans have predominantly heparan sulfate-GAG (HS-GAG) chains attached to the extracellular domain and in the case of Sdc1 and Sdc3, additionally chondroitin sulfate GAG chains (Deepa *et al.*, 2004).

The extracellular domain of all syndecans contains a proteinase-sensitive dibasic lysine-arginine-site (KR in Sdc2 and Sdc4) or arginine-lysine-site (RK in Sdc1, Sdc2 and Sdc3) adjacent to the transmembrane domain. This site was proposed to be a cleavage site for plasmin and thrombin. In 1989, the shedding of Sdc1 at this site was described by (Saunders *et al.*, 1989). Also, thrombin cleaved the Sdc4 at the Lys¹¹⁴-Arg¹¹⁵ link (Schmidt *et al.*, 2005). There are currently no data on the plasmin-dependent and thrombin-dependent shedding of Sdc2 and Sdc3, but it is likely that both can be shed at the respective sites. Furthermore, as discussed later in this review, several different shedding sites are located in the extracellular part of the protein.



Figure 1

Sequence alignment of all syndecan family members. The alignment indicates the different regions for each syndecan, including signal peptide (green), extracellular domain (yellow), transmembrane domain (blue) and intracellular domain (red). Furthermore, the homodimerization domain within the transmembrane region is marked in grey. Black bars indicate the C1, C2 and V region within the intracellular domain (Uniprot UniProt website at <http://www.uniprot.org/>).

Syndecan expression

Syndecans are involved in growth control, cell spreading, cellular recognition, cellular adhesion and signal transduction (Couchman, 2010; Choi *et al.*, 2011). Each syndecan has a tissue-specific and developmentally regulated pattern of expression (Kim *et al.*, 1994). For example, during murine development, Sdc1 is expressed first in the ectoderm and later on in mesodermal cells (Sutherland *et al.*, 1991). In mature tissue, epithelial cells permanently express Sdc1 (Kim *et al.*, 1994). The expression of Sdc3 is found during skeletal and neural development, where it is supposed to complement the function of Sdc1 (Bernfield *et al.*, 1993; Kim *et al.*, 1994). Sdc2 and Sdc4 are also expressed during mouse embryogenesis, specifically during endochondral ossification (David, 1993; Bertrand *et al.*, 2013). In contrast to the other syndecans, Sdc4 is expressed ubiquitously (Kim *et al.*, 1994).

As mentioned before, syndecans are mediators of various cellular functions. One explanation for how syndecans can fulfil all these functions might be the differential regulation of their expression during development and disease. For example, Sdc4 exhibits a major function in regulating cell matrix remodelling under inflammatory conditions, such as wound healing, fracture healing and osteoarthritis (Echtermeyer *et al.*, 2001; Echtermeyer *et al.*, 2009; Bertrand *et al.*, 2013). To exert this function, the expression of Sdc4 is regulated in an NF κ B-dependent manner, thereby explaining the increased expression under inflammatory conditions (Zhang *et al.*, 1999; Wang *et al.*, 2014). Apart from the specific regulation of expression, syndecans are able to initiate downstream signalling cascades via the C1 region, located just beneath the membrane, which is thought to interact with the cell cytoskeleton and cellular Src kinase proteins (Kinnunen *et al.*, 1998). Also, the C2 region, which contains a PDZ1 or PDZ2 domain, binds adaptor proteins, and the interaction mediates vesicular trafficking and exosome biogenesis (Gao *et al.*, 2000; Baietti *et al.*, 2012). The V region is thought to determine the role of syndecans in downstream signalling processes (Afratis *et al.*, 2017).

Beside this direct syndecan-dependent activation of signalling cascades, syndecans interact via their HS-GAG chains with a variety of ligands such as growth factors, cytokines, proteinases, adhesion receptors, extracellular matrix (ECM) components and morphogens (Pap and Bertrand, 2013). These HS-protein interactions are evolutionarily conserved and strongly HS-sequence and especially sequence modification specific. Various enzymes are needed for the maturation of HS-GAG chains, including multiple glycosyltransferases, sulfotransferases and an epimerase. It is known that many different cell types produce HS chains with several post-translational modifications, which determine the activation of downstream signalling cascades (Gesteira *et al.*, 2011; Shah *et al.*, 2011; Mortier *et al.*, 2012). This massive influence of HS-GAG chain modifications on signal transduction is explained by the fact that these modifications modulate the binding capacity of morphogens and chemokines, as, for example, the 6-O-sulfation of HS-GAG chains seems to be necessary for activation of FGF and Wnt signalling (Dhoot *et al.*, 2001; Wang *et al.*, 2004).

Interestingly, mutations in most of these HS-GAG modifying enzymes are associated with different diseases, including various malfunctions during skeletal development (Koziel

et al., 2004; Kluppel *et al.*, 2005; Otsuki *et al.*, 2008; Otsuki *et al.*, 2010; Otsuki *et al.*, 2017) and neuronal network formation (Rhiner and Hengartner, 2006).

Upon binding of different morphogens to the GAG chains, syndecans on the one hand interact with the respective receptor at the cell surface. On the other hand, it has been shown that syndecans can be shed from the cell surface to build morphogen gradients throughout the ECM, making the shedding a relevant process in syndecan-dependent signalling pathways.

Shedding

It has been known for many years that syndecans link the cytoskeleton to the ECM (Rapraeger *et al.*, 1986). One of the first publications about syndecan shedding described this process as an attempt of cells to release themselves from this interaction with the ECM by a proteolytic cleavage (Jalkanen *et al.*, 1987). Today, it is known that under physiological conditions, the ectodomains of syndecans are constitutively shed to a small degree. This shedding rate can be substantially increased in response to external stimuli (Kim *et al.*, 1994; Manon-Jensen *et al.*, 2010). This review gives a broad overview about different pathways and mechanisms activated and/or modulated by the shed syndecans. There are certainly more sheddases and downstream activated pathways, which are not mentioned in this section, which are detailed in shedding specific reviews (Manon-Jensen *et al.*, 2010; Nam and Park, 2012).

Different sheddases are able to cleave syndecans on the extracellular side, releasing a soluble syndecan consisting of the extracellular domain and the attached GAG chains (Brule *et al.*, 2006; Pruessmeyer *et al.*, 2010). These soluble syndecans may function as paracrine or autocrine effectors, or function as decoy receptors by competing for the same ligands as their cell bound counterparts (Kainulainen *et al.*, 1998) (Figure 2A and B). These cleaved fragments contain intact HS-GAG chains that retain biological activity similar to that of their parent molecule. These fragments still have the ability to down-regulate signal transduction by competing with the membrane-bound syndecans for extracellular ligand binding and sequestering the HS binding factors in ECM (Hayashida *et al.*, 2008) (Figure 2A).

Soluble ectodomains of syndecans, however, do not only function as competitive inhibitors but can also work as agonists. For example, the ectodomain of Sdc1 binds to FGF-2 more efficiently than the cell surface bound Sdc1 and inhibits its mitogenicity (Su *et al.*, 2007). Upon degradation of the GAG chains attached to the soluble Sdc1 ectodomain by heparanase present in the wound fluids, FGF-2 is activated to enhance wound repair (Kato *et al.*, 1998; Yang *et al.*, 2002; Mahtouk *et al.*, 2007). Hence, syndecans have diverse functions both as membrane bound and soluble forms. Therefore, soluble syndecans can also help form morphogen gradients across tissues that influence cell behaviour, for example, migration in tissue repair (Li *et al.*, 2002; Manon-Jensen *et al.*, 2010) (Figure 2B).

Furthermore, heparanase, an endo- β -D-glucuronidase, plays a role in the shedding of syndecans. This fact is counter-intuitive, as heparanase is known only to cleave the HS-




Figure 2

Schematic representation of different effects of shed syndecans on signalling cascades. The diagram depicts the various signalling influences of soluble syndecans on downstream signalling cascades that are discussed in the text. (A) Decoy receptor; (B) morphogen gradient; (C) heparanase-induced shedding; (D) receptor activation by shed Sdc1; (E) C-terminal fragment signalling; (F) shed syndecan in the nucleus.

GAG chains of proteoglycans but not the core protein of syndecans (Pikas *et al.*, 1998). The enhanced shedding of syndecans activated by heparanase is mediated indirectly, via the up-regulation of MMP9 and induction of ERK signalling (Purushothaman *et al.*, 2008) (Figure 2C). It has been shown that the 2-O-sulfated HS-GAG chains of Sdc1 inhibit neutrophil-dependent cathelicidin secretion, thereby promoting *Staphylococcus aureus* infection (Hayashida *et al.*, 2015). These data indicate that the main effect of heparanase on syndecan shedding is the activation of inflammatory signalling cascades, inducing the production of sheddases.

Soluble syndecans can also help form morphogen gradients due to the factors attached to the HS-GAG chains. Furthermore, the cleaved HS-GAG chains themselves can induce cellular responses. Interestingly, shedding of Sdc1 exposes a cryptic domain on the soluble core protein that contains binding sites for VLA4 and VEGF receptor 2 (VEGFR2). The shed Sdc1 activates VEGFR2 and stimulates thereby tumour cell invasion (Jung *et al.*, 2016) (Figure 2D).

During the shedding of the external part of syndecan, also, a C-terminal membrane-bound fragment is generated (Fitzgerald *et al.*, 2000). These C-terminal syndecan fragments are further cleaved at the transmembrane region by presenilin-dependent γ secretase upon ectodomain shedding (Schulz *et al.*, 2003). The C-terminal Sdc2 fragment up-regulates MMP7 expression via the protein kinase C γ -mediated focal adhesion kinase (FAK)/ERK signalling pathway in colon cancer, thereby up-regulating its own shedding (Jang *et al.*, 2017) (Figure 2E).

Interestingly, the shed syndecan fragment can be taken up by cells, as it has been shown that shed Sdc1 translocated to the nucleus of cells delivering growth factors and inhibiting histone acetylation (Stewart *et al.*, 2015) (Figure 2F).

Syndecan ectodomain shedding is mediated by various MMPs, such as MMP2, MMP7 and MMP9 (Schlondorff and Blobel, 1999; Arribas and Borroto, 2002). Furthermore, plasmin and thrombin have been shown to function as sheddases for syndecans (Schmidt *et al.*, 2005; Wang *et al.*, 2005).

Shedding of Sdc1 and Sdc4 is accelerated by activation of thrombin and the EGF. This shows that proteases and growth factors, which are active during wound repair, can accelerate syndecan shedding from cell surfaces (Subramanian *et al.*, 1997). Interestingly, MMPs cleave syndecans at the juxtamembrane site in a process that is usually accelerated during diseased conditions (Manon-Jensen *et al.*, 2010). The disintegrin and metalloproteinases (ADAMs), however, cleave Sdc4 near the N-terminal tip of the first HS-GAG chain attachment site (Gao *et al.*, 2004; Rodriguez-Manzanique *et al.*, 2009) (Figure 3).

Sdc1 contains the general consensus motif for cleavage by MMP7, MMP9 and MMP14, and *in vitro* and *in vivo* evidence of shedding has been published (Li *et al.*, 2002; Endo *et al.*, 2003). Chen *et al.* (2009) showed that epithelial injury induced Sdc1 shedding from the epithelium of wild-type mice but not from the epithelium of MMP7 knockout mice, indicating an essential role for MMP7 in the shedding process. A very recent study showed that MMP14 sheds Sdc1 during liver fibrosis, where the soluble Sdc1 interferes with TGF- β 1 signalling and thereby up-regulates its own sheddase (Regos *et al.*, 2018). The gelatinases MMP2 and MMP9 have been shown to shed Sdc1, Sdc2 and Sdc4 (Brule *et al.*, 2006; Fears *et al.*, 2006). Controversial data have been published on the involvement of TNF- α -converting enzyme, ADAM-17 (TACE), in the shedding of syndecans. Fitzgerald found that ectodomain shedding of Sdc1 and Sdc4 is TACE independent (Fitzgerald *et al.*, 2000). Later, it was found that the shedding of Sdc1 and Sdc4 is stimulated by the recombinant TACE catalytic domain (Pruessmeyer *et al.*, 2010). Sdc3 shedding has been reported in Schwann cells obtained from the sciatic nerves of 2- to 4-day-old rats (Asundi *et al.*, 2003). As the shedding process was reduced in cells treated with an MMP inhibitor, the involvement of MMPs in mediating Sdc3 shedding is very likely (Asundi *et al.*, 2003) (Figure 3).

These results indicate that syndecans can be the substrate of more than one sheddase, suggesting that different sheddases act in a tissue-specific manner. The different functions of the various shed fragments and their attached factors are still not fully understood, but it has become clear that shed syndecans influence signalling cascades in several different ways. As shedding of syndecans is specifically regulated under disease conditions, soluble syndecan ectodomains are used as biomarker for various diseases.

Shed syndecans as biomarkers for different diseases

Syndecan shedding has been shown to regulate many pathophysiological processes, such as inflammation, tissue repair and cancer cell proliferation (Maeda *et al.*, 2004). Tissue injury is accompanied by cellular stress, accumulation of leukocyte-derived proteases (thrombin, plasmin, elastase, etc.) and release of growth factors, each of which may accelerate syndecan shedding (Subramanian *et al.*, 1997). For this reason, shed syndecan ectodomains are found in inflammatory fluids, where they are thought to maintain proteolytic and growth factor balance (Subramanian *et al.*, 1997), as well as mediating inflammation (Fitzgerald *et al.*, 2000). A detailed list of the different diseases for which shed Sdc1, Sdc2, Sdc3 and Sdc4 have been proposed as biomarkers is given in Table 1. We provide here some typical studies on shed syndecans as biomarkers for various diseases.

Most publications on soluble syndecans as biomarkers focus on soluble Sdc1. For example, Sdc1 ectodomains are elevated in blood of patients with sepsis (Nelson *et al.*, 2008; Steppan *et al.*, 2011), ischaemia-reperfusion injury (Rehm *et al.*, 2007), graft-versus-host disease (Seidel *et al.*, 2003) and various cancers (Joensuu *et al.*, 2002; Yang *et al.*, 2002). Furthermore, studies in mice have shown that the




Figure 3

Overview of the proteinase cleavage sites on the syndecan ectodomains. The validated shedding sites for Sdc1 and Sdc4 are indicated at the respective amino acid number and sheddases according to Manon-Jensen *et al.* (2010) (shown in black). The predicted shedding sites for Sdc2 and Sdc3 include only the prediction for MMP2, MMP3 and MMP14 and thrombin. The shedding sites for Sdc2 and Sdc3 are generated using a cleavage site prediction tool (shown in grey). Again, the amino acid is indicated with the predicted sheddase. HS-GAG chains are shown in blue colour and chondroitin sulfate GAG chains are shown in orange.

Table 1

List of soluble syndecans as biomarkers for various diseases including the respective reference

Sdc1			
Disease	Sample	Regulation	References
Sepsis	Arterial plasma, serum	Significantly elevated levels, correlation with cardiovascular Sequential Organ Failure Assessment score	Nelson <i>et al.</i> (2008); Steppan <i>et al.</i> (2011)
Sepsis survival after major abdominal surgery	Plasma	Patients with post-operative sepsis showed increased levels; levels associated with survival after sepsis	Holzmann <i>et al.</i> (2018)
Acute traumatic endotheliopathy in isolated severe brain injury	Plasma	Sdc1 levels above 30.5 ng·mL ⁻¹ indicate patients with traumatic brain injury-associated coagulopathy	Albert <i>et al.</i> (2018)
Trauma patients	Serum/plasma	High levels of Sdc1 are associated with inflammation, coagulopathy and increased mortality, a syndecan-1 level ≥ 40 ng·mL ⁻¹ identified patients with worse outcome	Johansson <i>et al.</i> (2011); Johansson <i>et al.</i> (2012); Gonzalez Rodriguez <i>et al.</i> (2017)
Microvascular glycocalyx degradation	Plasma	Syndecan-1 correlates with glycocalyx thickness and permeability changes	Torres Filho <i>et al.</i> (2016)
KD	Serum	Sdc1 levels may indicate endothelial damage and inflammation KD	Luo <i>et al.</i> (2018a)
Pulmonary embolism	Blood	Increased levels of Sdc1 in high risk pulmonary embolism patients	Lehnert <i>et al.</i> (2017)
Multicentric Castleman's disease	Bronchoalveolar lavage fluid	Marked elevation of soluble Sdc1	Hasegawa <i>et al.</i> (2007)
Heart failure	Plasma	Syndecan-1 correlates with fibrosis biomarkers	Tromp <i>et al.</i> (2014)
Ventricular remodelling after myocardial infarction	Serum	Increased levels of soluble Sdc1	Lei <i>et al.</i> (2012)
Takotsubo cardiomyopathy	Blood	Sdc1 is significantly increased in the acute stage of TCC	Nguyen <i>et al.</i> (2017)
AKI	Blood	Prognostic marker to assess the risk of AKI	Liborio <i>et al.</i> (2015); Neves <i>et al.</i> (2015); de Melo Bezerra Cavalcante <i>et al.</i> (2016)
Chronic kidney disease	Plasma	Plasma levels were increased compared to the healthy control group	Padberg <i>et al.</i> (2014)
Ischaemia–reperfusion injury	Arterial blood	Elevated levels of Sdc1	Rehm <i>et al.</i> (2007)
DIC	Serum/plasma	Correlates with increased levels of Sdc1, predicts DIC in patients with sepsis	Ikeda <i>et al.</i> (2018)
Hypocoagulation	Serum/plasma	Increased levels are associated with hypocoagulation in patients with sepsis	Ostrowski <i>et al.</i> (2015)
GVHD	Serum	Sdc1 levels elevated in patients who developed acute GVHD after allogeneic stem cell transplantation	Seidel <i>et al.</i> (2003)
Crohn's disease	Serum	Higher Sdc1 levels compared to normal population	Zhang <i>et al.</i> (2013); Cekic <i>et al.</i> (2015)
Small bowel damage in children with CD	Serum	Elevated levels of Sdc1 in children with CD, correlation of Sdc1 levels and mucosal damage	Yablecovitch <i>et al.</i> (2017)
SLE	Serum	Higher levels in SLE patients with nephritis compared to RA patients and healthy control group, possible marker for active SLE	Minowa <i>et al.</i> (2011); Kim <i>et al.</i> (2015); Mosaad <i>et al.</i> (2017)
Liver fibrosis stage in patients with hepatitis C	Serum	Suggested as non-invasive marker to predict liver fibrosis stage	Zvibel <i>et al.</i> (2009)
Hantavirus infection	Plasma	Sdc1 was associated with disease severity (as well as levels of thrombocytes, albumin, IGFBP-1, decreased blood pressure)	Connolly-Andersen <i>et al.</i> (2014)
Type I diabetes mellitus	Serum	Sdc1 is upregulated	Svennevig <i>et al.</i> (2006)

continues

Table 1

(Continued)

Sdc1			
Disease	Sample	Regulation	References
PE	Serum/plasma	Statistical differences in serum between PE and normal pregnancy, Sdc1 in plasma is significantly lower before the onset of PE	Gandley <i>et al.</i> (2016); Alici Davutoglu <i>et al.</i> (2018)
HELLP syndrome	Serum	Sdc1 levels increase in normal pregnancy but even higher in women with HELLP	Hofmann-Kiefer <i>et al.</i> (2013)
Rhegmatogenous retinal detachment	Subretinal fluid/vitreous fluid	Significant increase of Sdc1	Wang <i>et al.</i> (2008)
Systemic sclerosis	Serum	Significantly higher than in healthy control group	Wu <i>et al.</i> (2016)
Pleural malignancies	Pleural effusions	Sdc1 levels can distinguish malignant and benign disease	Mundt <i>et al.</i> (2014)
Lung cancer	Serum	High Sdc1 levels were associated with a poor survival rate	Joensuu <i>et al.</i> (2002); Anttonen <i>et al.</i> (2003)
Myeloma (multiple)	Serum	Possible prognostic marker	Dhodapkar <i>et al.</i> (1998); Seidel <i>et al.</i> (2000); Yang <i>et al.</i> (2002); Aref <i>et al.</i> (2003); Janosi <i>et al.</i> (2004); Lovell <i>et al.</i> (2005); Maisnar <i>et al.</i> (2006); Scudla <i>et al.</i> (2009); Kim <i>et al.</i> (2010)
Hepatocellular carcinoma	Serum	High levels in patients with hepatocellular carcinoma detected, high levels associated with greater risk of tumour recurrence and death	Metwaly <i>et al.</i> (2012); Nault <i>et al.</i> (2013)
(Metastatic) CRC	Serum	Baseline Sdc1 is suggested as prognostic marker for overall survival in metastatic CRC, Sdc1, among others, may be involved in tumour progression and can be used for prognosis of CRC patients	Jary <i>et al.</i> (2016); Mitselou <i>et al.</i> (2016)
PC	Serum	Significant higher Sdc1 levels in advanced cases of PC, independent factor of adverse overall and disease-specific survival	Szarvas <i>et al.</i> (2016)
Hodgkin's lymphoma	Serum	Serum levels are elevated but do not strongly correlate with other parameters, further evaluation is required	Vassilakopoulos <i>et al.</i> (2005)
Lymphocytic leukaemia	Plasma	Soluble Sdc1 in combination with beat2-M and Rai stage may replace testing for IgVH mutation status	Jilani <i>et al.</i> (2009)
Breast cancer	Serum	Positive correlation between soluble Sdc1 and tumour size	Malek-Hosseini <i>et al.</i> (2017)
Bladder cancer	Serum	Increased levels in sera of bladder cancer patients	Sanaee <i>et al.</i> (2015)
Sdc2			
Disease	Sample	Regulation	References
Colon cancer	Serum	Sdc2 detectable in majority of colon cancer patients, while all healthy patients were negative	Choi <i>et al.</i> (2015)
Keloid tissue	Tissue	Up-regulated in keloid tissue	Mukhopadhyay <i>et al.</i> (2010)
Sdc3			
Soluble Sdc3 has not been reported to be a suitable biomarker.			
Sdc4			
Disease	Sample	Regulation	References
Acute bacterial pneumonia	Serum	Up-regulation (at the onset, mild pneumonia compared to severe pneumonia)	Nikaïdo <i>et al.</i> (2015)

continues

Table 1

(Continued)

Sdc4			
Disease	Sample	Regulation	References
IIP	Serum	Down-regulation in acute exacerbation, patients with higher baseline have worse prognosis for IIP (prognostic predictor)	Sato <i>et al.</i> (2017)
Severe community acquired pneumonia	Serum	Patients with Sdc4 levels below 6.68 ng·mL ⁻¹ have a higher mortality (prognostic predictor)	Luo <i>et al.</i> (2018b)
Heart failure in patients with hypertension	Serum	Sdc4 levels above 2.3 ng·mL ⁻¹ (among others) are significant predictor for heart failure	Bielecka-Dabrowa <i>et al.</i> (2015)
Adverse LV remodelling in patients with dilated cardiomyopathy	Serum	Sdc4 correlated positively with LV diastolic/systolic diameters, might be useful in predicting LV remodelling	Takahashi <i>et al.</i> (2011); Bielecka-Dabrowa <i>et al.</i> (2013)
Atopic dermatitis	Serum	Levels upregulated compared to control group and correlate with disease severity, eczema area, severity index and itch visual analogue scale scores	Nakao <i>et al.</i> (2016)
Cardiovascular mortality in HD patients	Serum	Sdc4 levels are increased in HD patients, levels correlate with echocardiographic parameters (predictor for cardiovascular mortality)	Jaroszynski <i>et al.</i> (2016)

AKI, acute kidney injury; CD, celiac disease; CRC, colorectal cancer; DIC, disseminated intravascular coagulation; KD, Kawasaki disease; GVHD, Graft-versus-host disease; HELLIP, Haemolysis, elevated liver enzymes and low platelets; HD, haemodialysis; IIP, idiopathic intestinal pneumonia; PC, prostate cancer; PE, preeclampsia; RA, rheumatoid arthritis; SLE, Systemic lupus erythematosus; TCC, terminal complement complex.

inflammatory response to toxins, chemicals, allergens and pathogens is dysregulated in the absence of Sdc1 or when its shedding is inhibited (Kainulainen *et al.*, 1998; Kato *et al.*, 1998), suggesting that Sdc1 shedding is activated to ensure adequate tissue response to inflammation. Consequently, Sdc1 has been proposed to be a biomarker for sepsis survival after major abdominal surgery, as well as for acute traumatic endotheliopathy in isolated severe brain injury, and for small bowel mucosal damage in children with celiac disease (Yablecovitch *et al.*, 2017; Albert *et al.*, 2018; Holzmann *et al.*, 2018).

The activation of Sdc2 shedding has been described for cancer cells. In particular, the MMP7-induced shedding of Sdc2 was detected in colon cancer cells *in vitro* (Choi *et al.*, 2011). Patients with advanced colon cancer exhibited significantly higher Sdc2 serum levels compared to a healthy control group, which was mainly negative for Sdc2 serum levels (Choi *et al.*, 2015). Furthermore, Sdc2 and FGF-2 were overexpressed in keloid tissue. The authors suggested that both proteins interact with each other, resulting in shedding of Sdc2 and that shed Sdc2 might be involved in the keloidic phenotype (Mukhopadhyay *et al.*, 2010).

Besides its role in cancer, shed Sdc2 has been linked to angiogenesis, as its expression is increased during endothelial cell angiogenic processes (Fears *et al.*, 2006). Shed Sdc2 regulated angiogenesis by inhibiting endothelial cell migration in human and rodent models and thereby reduced tumour growth (De Rossi *et al.*, 2014).

Soluble Sdc4 in serum is mainly associated with pneumonia and heart failure. In patients with mild pneumonia, Sdc4 was increased in comparison to patients with severe pneumonia. Interestingly, a short-term antibiotic therapy further increased Sdc4 levels, leading the authors to the suggestion that Sdc4 might have an anti-inflammatory function (Nikaido *et al.*, 2015). The same research group

showed that Sdc4 levels were increased in patients with idiopathic interstitial pneumonia. Again, the authors propose that baseline serum Sdc4 levels were indicative for the prognosis, showing that higher serum levels of Sdc4 were associated with a worse prognosis than lower baseline levels (Sato *et al.*, 2017). Sdc4 serum levels were also associated with severe community-acquired pneumonia, and these increased serum levels were linked to a higher mortality rate (Luo *et al.*, 2018b).

Furthermore, high serum Sdc4 levels were found to be a significant predictor of heart failure in patients with hypertension (Bielecka-Dabrowa *et al.*, 2015), and Sdc4 was also proposed as a suitable biomarker for the adverse left ventricular (LV) remodelling in patients with dilated cardiomyopathy (Bielecka-Dabrowa *et al.*, 2013). This finding was corroborated in another study in which serum Sdc4 levels were proposed to be a biomarker for LV remodelling in heart failure (Takahashi *et al.*, 2011). Serum Sdc4 levels were also increased in haemodialysis patients and correlated with geometrical echocardiographic parameters. This study suggested Sdc4 as a predictor for cardiovascular mortality in haemodialysis patients (Jaroszynski *et al.*, 2016). Apart from cardiovascular diseases and pneumonia, increased Sdc4 serum levels have also been reported in patients with atopic dermatitis. In this study, Sdc4 levels correlated with the disease severity as well as eczema area, the severity index and visual analogue scale scores for itch (Nakao *et al.*, 2016).

In particular, serum levels of soluble Sdc1 and Sdc4 have been shown to be associated with various diseases. Therefore, it will be a challenge to differentiate the cause of elevated syndecan levels in different patients to decide to which disease they might relate. More distinct analyses of the shedding site and/or modification of the HS-GAG chains will improve the sensitivity of the suggested biomarker for the respective disease.

inhibit myeloma cell viability *in vitro* and *in vivo* in a mouse model of breast cancer (Pumphrey *et al.*, 2002). Not only soluble Sdc1 has been shown to exhibit therapeutic potential, there is also evidence for a beneficial effect of shed Sdc2. Shed Sdc2 inhibits angiogenesis by inhibiting endothelial cell migration and thereby reduces tumour growth (De Rossi *et al.*, 2014). This finding gives rise to a novel therapeutic strategy to target pathologies that are characterized by new blood vessel formation, like different cancers, infectious diseases and autoimmune disorders.

These studies highlight the anticancer and antiviral activities of GAG chain-containing proteins and provide the foundation for future development of synthetic proteoglycans as novel therapeutic agents.

Interestingly, also the overexpression of the C-terminal fragment of Sdc1 has been shown to suppress migration and invasion of tumour cells. This inhibitory effect, however, was only seen in cells expressing endogenous Sdc1 but not in Sdc1 knockout cells. The C-terminal Sdc1 fragment suppressed tumour cell migration and increased basal phosphorylation of Src and FAK. The authors explain the observed effects with an antagonizing mechanism of the C-terminal fragment for the Sdc1-dependent tumour cell migration *in vitro* and *in vivo* by dysregulating pro-adhesive signalling pathways (Pasqualon *et al.*, 2015).

There is also evidence that the blockade of Sdc4 using a blocking antibody might have a positive effect in preventing cartilage destruction in a mouse model of osteoarthritis (Echtermeyer *et al.*, 2009). The authors describe that Sdc4-mediated cartilage destruction in osteoarthritis is mediated by binding of the aggrecanase (ADAMTS-5) to the side chains of Sdc4, thereby fixing ADAMTS-5 at the cell surface. The activation of ADAMTS-5 is mediated by MMP3 expression, which is regulated in an IL-1-dependent manner by Sdc4, as Sdc4 regulates the sensitivity of chondrocytes to IL-1 signalling (Echtermeyer *et al.*, 2009).

Figure 4 summarizes the current therapeutic strategies involving modification of Sdc-dependent signalling pathways. Four different approaches can be differentiated. The first approach is based on the inhibition of HS side chain cleavage and thereby inhibits the HS-fragment-induced activation of inflammatory signalling cascades and Sdc-shedding (Figure 4A). The second strategy is based on the anti-inflammatory effect of low MW heparin, although the exact mechanism of the anti-inflammatory effect is not known (Figure 4B). The third mechanism is based on the blockade of Sdc4 signalling and inhibition of IL-1-dependent inflammatory signals. Again, the exact mechanism of this blockade is not described (Figure 4C). The last approach is based on the use of soluble syndecans or their synthetic variants. There are several studies using either full length soluble syndecans or truncated forms or even synthetic variants with synthetic HS side chains attached (Figure 4D).

Conclusion

The current knowledge about syndecan shedding highlights the role of soluble syndecans in various diseases. However, the main function of soluble syndecans depends mainly on the presence of GAG chains, which are known to be

modified during ageing, disease and cell differentiation (Bassett *et al.*, 2006). At the same time, the respective sheddases are modulated depending on external stimuli, cell differentiation and inflammation. The current knowledge just elucidates parts of the highly complex temporal and spatial regulation of syndecan expression, regulation of morphogen binding and further shedding during ageing and diseases. This makes the usage of soluble syndecans as biomarkers difficult, especially as more than one stimulus might evoke shedding of the same syndecan, thereby reducing the specificity of the potential biomarker. There is clearly a therapeutic potential for soluble syndecans in different diseases; however, more insight in the role GAG chains and GAG chain modification is needed to fully understand the different roles and effects. Syndecan core proteins most likely serve mainly as the anchorage for these highly complex sugar chains, building the basis for disease regulated shedding.

Nomenclature of targets and ligands

Key protein targets and ligands in this article are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Harding *et al.*, 2018), and are permanently archived in the Concise Guide to PHARMACOLOGY 2017/18 (Alexander *et al.*, 2017a,b).

Acknowledgements

We would like to thank the DFG for funding (Emmy Noether BE4328/5-1).

Conflict of interest

The authors declare no conflicts of interest.

References

- Afratis NA, Nikitovic D, Multhaupt HA, Theocharis AD, Couchman JR, Karamanos NK (2017). Syndecans – key regulators of cell signaling and biological functions. *FEBS J* 284: 27–41.
- Albert V, Subramanian A, Agrawal D, Pati HP, Gupta SD, Mukhopadhyay AK (2018). Acute traumatic endotheliopathy in isolated severe brain injury and its impact on clinical outcome. *Med Sci (Basel)* 6: 5.
- Alexander SPH, Fabbro D, Kelly E, Marrion NV, Peters JA, Faccenda E *et al.* (2017a). The Concise Guide to PHARMACOLOGY 2017/18: Enzymes. *Br J Pharmacol* 174: S272–S359.
- Alexander SPH, Fabbro D, Kelly E, Marrion NV, Peters JA, Faccenda E *et al.* (2017b). The Concise Guide to PHARMACOLOGY 2017/18: Catalytic receptors. *Br J Pharmacol* 174: S225–S271.
- Alici Davutoglu E, Akkaya Firat A, Ozel A, Yilmaz N, Uzun I, Temel Yuksel I *et al.* (2018). Evaluation of maternal serum hypoxia inducible factor-1alpha, progranulin and syndecan-1 levels in pregnancies with early- and late-onset preeclampsia. *J Matern Fetal Neonatal Med* 31: 1976–1982.

- Anttonen A, Leppä S, Ruotsalainen T, Alftan H, Mattson K, Joensuu H (2003). Pretreatment serum syndecan-1 levels and outcome in small cell lung cancer patients treated with platinum-based chemotherapy. *Lung Cancer* 41: 171–177.
- Aref S, Goda T, El-Sherbiny M (2003). Syndecan-1 in multiple myeloma: relationship to conventional prognostic factors. *Hematology* 8: 221–228.
- Arribas J, Borroto A (2002). Protein ectodomain shedding. *Chem Rev* 102: 4627–4638.
- Asundi VK, Erdman R, Stahl RC, Carey DJ (2003). Matrix metalloproteinase-dependent shedding of syndecan-3, a transmembrane heparan sulfate proteoglycan, in Schwann cells. *J Neurosci Res* 73: 593–602.
- Baietti MF, Zhang Z, Mortier E, Melchior A, Degeest G, Geeraerts A *et al.* (2012). Syndecan-syntenin-ALIX regulates the biogenesis of exosomes. *Nat Cell Biol* 14: 677–685.
- Bassett JH, Swinhoe R, Chassande O, Samarut J, Williams GR (2006). Thyroid hormone regulates heparan sulfate proteoglycan expression in the growth plate. *Endocrinology* 147: 295–305.
- Beauvais DM, Jung O, Yang Y, Sanderson RD, Rapraeger AC (2016). Syndecan-1 (CD138) suppresses apoptosis in multiple myeloma by activating IGF1 receptor: prevention by Synstatin IGF1R inhibits tumor growth. *Cancer Res* 76: 4981–4993.
- Bernfield M, Hinkes MT, Gallo RL (1993). Developmental expression of the syndecans: possible function and regulation. *Dev Suppl*: 205–212.
- Bertrand J, Stange R, Hidding H, Echtermeyer F, Nalesso G, Godmann L *et al.* (2013). Syndecan 4 supports bone fracture repair, but not fetal skeletal development, in mice. *Arthritis Rheum* 65: 743–752.
- Bielecka-Dabrowa A, Gluba-Brzozka A, Michalska-Kasiczak M, Misztal M, Rysz J, Banach M (2015). The multi-biomarker approach for heart failure in patients with hypertension. *Int J Mol Sci* 16: 10715–10733.
- Bielecka-Dabrowa A, von Haehling S, Aronow WS, Ahmed MI, Rysz J, Banach M (2013). Heart failure biomarkers in patients with dilated cardiomyopathy. *Int J Cardiol* 168: 2404–2410.
- Bobardt MD, Chatterji U, Schaffer L, de Witte L, Gallay PA (2010). Syndecan-Fc hybrid molecule as a potent in vitro microbicidal anti-HIV-1 agent. *Antimicrob Agents Chemother* 54: 2753–2766.
- Brule S, Charnaux N, Sutton A, Ledoux D, Chaigneau T, Saffar L *et al.* (2006). The shedding of syndecan-4 and syndecan-1 from HeLa cells and human primary macrophages is accelerated by SDF-1/CXCL12 and mediated by the matrix metalloproteinase-9. *Glycobiology* 16: 488–501.
- Carey DJ (1997). Syndecans: multifunctional cell-surface co-receptors. *Biochem J* 327 (Pt 1): 1–16.
- Cekic C, Kirci A, Vatansever S, Aslan F, Yilmaz HE, Alper E *et al.* (2015). Serum syndecan-1 levels and its relationship to disease activity in patients with Crohn's disease. *Gastroenterol Res Pract* 2015: 850351.
- Chen P, Abacherli LE, Nadler ST, Wang Y, Li Q, Parks WC (2009). MMP7 shedding of syndecan-1 facilitates re-epithelialization by affecting $\alpha(2)\beta(1)$ integrin activation. *PLoS One* 4: e6565.
- Choi S, Choi Y, Jun E, Kim IS, Kim SE, Jung SA *et al.* (2015). Shed syndecan-2 enhances tumorigenic activities of colon cancer cells. *Oncotarget* 6: 3874–3886.
- Choi S, Lee E, Kwon S, Park H, Yi JY, Kim S *et al.* (2005). Transmembrane domain-induced oligomerization is crucial for the functions of syndecan-2 and syndecan-4. *J Biol Chem* 280: 42573–42579.
- Choi Y, Chung H, Jung H, Couchman JR, Oh ES (2011). Syndecans as cell surface receptors: unique structure equates with functional diversity. *Matrix Biol* 30: 93–99.
- Connolly-Andersen AM, Thunberg T, Ahlm C (2014). Endothelial activation and repair during hantavirus infection: association with disease outcome. *Open Forum Infect Dis* 1: ofu027.
- Couchman JR (2010). Transmembrane signaling proteoglycans. *Annu Rev Cell Dev Biol* 26: 89–114.
- David G (1993). Integral membrane heparan sulfate proteoglycans. *FASEB J* 7: 1023–1030.
- de Melo Bezerra Cavalcante CT, Castelo Branco KM, Pinto Junior VC, Meneses GC, de Oliveira Neves FM, de Souza NM *et al.* (2016). Syndecan-1 improves severe acute kidney injury prediction after pediatric cardiac surgery. *J Thorac Cardiovasc Surg* 152: 178–186 e172.
- De Rossi G, Evans AR, Kay E, Woodfin A, McKay TR, Nourshargh S *et al.* (2014). Shed syndecan-2 inhibits angiogenesis. *J Cell Sci* 127: 4788–4799.
- Deepa SS, Yamada S, Zako M, Goldberger O, Sugahara K (2004). Chondroitin sulfate chains on syndecan-1 and syndecan-4 from normal murine mammary gland epithelial cells are structurally and functionally distinct and cooperate with heparan sulfate chains to bind growth factors. A novel function to control binding of midkine, pleiotrophin, and basic fibroblast growth factor. *J Biol Chem* 279: 37368–37376.
- Dhodapkar MV, Abe E, Theus A, Lacy M, Langford JK, Barlogie B *et al.* (1998). Syndecan-1 is a multifunctional regulator of myeloma pathobiology: control of tumor cell survival, growth, and bone cell differentiation. *Blood* 91: 2679–2688.
- Dhoot GK, Gustafsson MK, Ai X, Sun W, Standiford DM, Emerson CP Jr (2001). Regulation of Wnt signaling and embryo patterning by an extracellular sulfatase. *Science* 293: 1663–1666.
- Echtermeyer F, Bertrand J, Dreier R, Meinecke I, Neugebauer K, Fuerst M *et al.* (2009). Syndecan-4 regulates ADAMTS-5 activation and cartilage breakdown in osteoarthritis. *Nat Med* 15: 1072–1076.
- Echtermeyer F, Streit M, Wilcox-Adelman S, Saoncella S, Denhez F, Detmar M *et al.* (2001). Delayed wound repair and impaired angiogenesis in mice lacking syndecan-4. *J Clin Invest* 107: R9–R14.
- Elenius K, Jalkanen M (1994). Function of the syndecans—a family of cell surface proteoglycans. *J Cell Sci* 107 (Pt 11): 2975–2982.
- Endo K, Takino T, Miyamori H, Kinsen H, Yoshizaki T, Furukawa M *et al.* (2003). Cleavage of syndecan-1 by membrane type matrix metalloproteinase-1 stimulates cell migration. *J Biol Chem* 278: 40764–40770.
- Fan L, Wu Q, Xing X, Liu Y, Shao Z (2011). Targeted silencing of heparanase gene by small interfering RNA inhibits invasiveness and metastasis of osteosarcoma cells. *J Huazhong Univ Sci Technolog Med Sci* 31: 348–352.
- Fears CY, Gladson CL, Woods A (2006). Syndecan-2 is expressed in the microvasculature of gliomas and regulates angiogenic processes in microvascular endothelial cells. *J Biol Chem* 281: 14533–14536.
- Fitzgerald ML, Wang Z, Park PW, Murphy G, Bernfield M (2000). Shedding of syndecan-1 and -4 ectodomains is regulated by multiple signaling pathways and mediated by a TIMP-3-sensitive metalloproteinase. *J Cell Biol* 148: 811–824.

- Gandley RE, Althouse A, Jeyabalan A, Bregand-White JM, McGonigal S, Myerski AC *et al.* (2016). Low soluble syndecan-1 precedes preeclampsia. *PLoS One* 11: e0157608.
- Gao G, Plaas A, Thompson VP, Jin S, Zuo F, Sandy JD (2004). ADAMTS4 (aggrecanase-1) activation on the cell surface involves C-terminal cleavage by glycosylphosphatidyl inositol-anchored membrane type 4-matrix metalloproteinase and binding of the activated proteinase to chondroitin sulfate and heparan sulfate on syndecan-1. *J Biol Chem* 279: 10042–10051.
- Gao Y, Li M, Chen W, Simons M (2000). Synectin, syndecan-4 cytoplasmic domain binding PDZ protein, inhibits cell migration. *J Cell Physiol* 184: 373–379.
- Gesteira TF, Coulson-Thomas VJ, Taunay-Rodrigues A, Oliveira V, Thacker BE, Juliano MA *et al.* (2011). Inhibitory peptides of the sulfotransferase domain of the heparan sulfate enzyme, N-deacetylase-N-sulfotransferase-1. *J Biol Chem* 286: 5338–5346.
- Gonzalez Rodriguez E, Ostrowski SR, Cardenas JC, Baer LA, Tomasek JS, Henriksen HH *et al.* (2017). Syndecan-1: a quantitative marker for the endotheliopathy of trauma. *J Am Coll Surg* 225: 419–427.
- Harding SD, Sharman JL, Faccenda E, Southan C, Pawson AJ, Ireland S *et al.* (2018). The IUPHAR/BPS Guide to PHARMACOLOGY in 2018: updates and expansion to encompass the new guide to IMMUNOPHARMACOLOGY. *Nucl Acids Res* 46: D1091–D1106.
- Hasegawa M, Betsuyaku T, Yoshida N, Nasuhara Y, Kinoshita I, Ohta S *et al.* (2007). Increase in soluble CD138 in bronchoalveolar lavage fluid of multicentric Castleman's disease. *Respirology* 12: 140–143.
- Hayashida A, Amano S, Gallo RL, Linhardt RJ, Liu J, Park PW (2015). 2-O-Sulfated domains in syndecan-1 heparan sulfate inhibit neutrophil cathelicidin and promote *Staphylococcus aureus* corneal infection. *J Biol Chem* 290: 16157–16167.
- Hayashida K, Chen Y, Bartlett AH, Park PW (2008). Syndecan-1 is an in vivo suppressor of Gram-positive toxic shock. *J Biol Chem* 283: 19895–19903.
- Hofmann-Kiefer KF, Knabl J, Martinoff N, Schiessl B, Conzen P, Rehm M *et al.* (2013). Increased serum concentrations of circulating glycocalyx components in HELLP syndrome compared to healthy pregnancy: an observational study. *Reprod Sci* 20: 318–325.
- Holzmann MS, Winkler MS, Strunden MS, Izbicki JR, Schoen G, Greiwe G *et al.* (2018). Syndecan-1 as a biomarker for sepsis survival after major abdominal surgery. *Biomark Med* 12: 119–127.
- Ikeda M, Matsumoto H, Ogura H, Hirose T, Shimizu K, Yamamoto K *et al.* (2018). Circulating syndecan-1 predicts the development of disseminated intravascular coagulation in patients with sepsis. *J Crit Care* 43: 48–53.
- Jalkanen M, Rapraeger A, Saunders S, Bernfield M (1987). Cell surface proteoglycan of mouse mammary epithelial cells is shed by cleavage of its matrix-binding ectodomain from its membrane-associated domain. *J Cell Biol* 105: 3087–3096.
- Jang B, Jung H, Choi S, Lee YH, Lee ST, Oh ES (2017). Syndecan-2 cytoplasmic domain up-regulates matrix metalloproteinase-7 expression via the protein kinase C γ -mediated FAK/ERK signaling pathway in colon cancer. *J Biol Chem* 292: 16321–16332.
- Janosi J, Sebestyen A, Mikala G, Nemeth J, Kiss Z, Valyi-Nagy I (2004). Soluble syndecan-1 levels in different plasma cell dyscrasias and in different stages of multiple myeloma. *Haematologica* 89: 370–371.
- Jaroszynski AJ, Jaroszynska A, Przywara S, Zaborowski T, Ksiazek A, Dabrowski W (2016). Syndecan-4 is an independent predictor of all-cause as well as cardiovascular mortality in hemodialysis patients. *PLoS One* 11: e0163532.
- Jary M, Lecomte T, Bouche O, Kim S, Dobi E, Queiroz L *et al.* (2016). Prognostic value of baseline seric syndecan-1 in initially unresectable metastatic colorectal cancer patients: a simple biological score. *Int J Cancer* 139: 2325–2335.
- Jilani I, Wei C, Bekele BN, Zhang ZJ, Keating M, Wierda W *et al.* (2009). Soluble syndecan-1 (sCD138) as a prognostic factor independent of mutation status in patients with chronic lymphocytic leukemia. *Int J Lab Hematol* 31: 97–105.
- Joensuu H, Anttonen A, Eriksson M, Makitaro R, Alfthan H, Kinnula V *et al.* (2002). Soluble syndecan-1 and serum basic fibroblast growth factor are new prognostic factors in lung cancer. *Cancer Res* 62: 5210–5217.
- Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR (2011). A high admission syndecan-1 level, a marker of endothelial glycocalyx degradation, is associated with inflammation, protein C depletion, fibrinolysis, and increased mortality in trauma patients. *Ann Surg* 254: 194–200.
- Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR (2012). High circulating adrenaline levels at admission predict increased mortality after trauma. *J Trauma Acute Care Surg* 72: 428–436.
- Jung O, Trapp-Stamborski V, Purushothaman A, Jin H, Wang H, Sanderson RD *et al.* (2016). Heparanase-induced shedding of syndecan-1/CD138 in myeloma and endothelial cells activates VEGFR2 and an invasive phenotype: prevention by novel synstatins. *Oncogene* 35: e202.
- Kainulainen V, Wang H, Schick C, Bernfield M (1998). Syndecans, heparan sulfate proteoglycans, maintain the proteolytic balance of acute wound fluids. *J Biol Chem* 273: 11563–11569.
- Kato M, Wang H, Kainulainen V, Fitzgerald ML, Ledbetter S, Ornitz DM *et al.* (1998). Physiological degradation converts the soluble syndecan-1 ectodomain from an inhibitor to a potent activator of FGF-2. *Nat Med* 4: 691–697.
- Kim CW, Goldberger OA, Gallo RL, Bernfield M (1994). Members of the syndecan family of heparan sulfate proteoglycans are expressed in distinct cell-, tissue-, and development-specific patterns. *Mol Biol Cell* 5: 797–805.
- Kim JM, Lee JA, Cho IS, Ihm CH (2010). Soluble syndecan-1 at diagnosis and during follow up of multiple myeloma: a single institution study. *Korean J Hematol* 45: 115–119.
- Kim KJ, Kim JY, Baek IW, Kim WU, Cho CS (2015). Elevated serum levels of syndecan-1 are associated with renal involvement in patients with systemic lupus erythematosus. *J Rheumatol* 42: 202–209.
- Kinnunen T, Kaksonen M, Saarinen J, Kalkkinen N, Peng HB, Rauvala H (1998). Cortactin-Src kinase signaling pathway is involved in N-syndecan-dependent neurite outgrowth. *J Biol Chem* 273: 10702–10708.
- Kluppel M, Wight TN, Chan C, Hinek A, Wrana JL (2005). Maintenance of chondroitin sulfation balance by chondroitin-4-sulfotransferase 1 is required for chondrocyte development and growth factor signaling during cartilage morphogenesis. *Development* 132: 3989–4003.
- Koziel L, Kunath M, Kelly OG, Vortkamp A (2004). Ext1-dependent heparan sulfate regulates the range of Ihh signaling during endochondral ossification. *Dev Cell* 6: 801–813.
- Lehnert P, Johansson PI, Ostrowski SR, Moller CH, Bang LE, Olsen PS *et al.* (2017). Coagulopathy in patients with acute pulmonary embolism: a pilot study of whole blood coagulation and markers of endothelial damage. *Scand J Clin Lab Invest* 77: 19–26.

- Lei J, Xue SN, Wu W, Zhou SX, Zhang YL, Yuan GY *et al.* (2012). Increased level of soluble syndecan-1 in serum correlates with myocardial expression in a rat model of myocardial infarction. *Mol Cell Biochem* 359: 177–182.
- Li Q, Park PW, Wilson CL, Parks WC (2002). Matrilysin shedding of syndecan-1 regulates chemokine mobilization and transepithelial efflux of neutrophils in acute lung injury. *Cell* 111: 635–646.
- Liborio AB, Braz MB, Seguro AC, Meneses GC, Neves FM, Pedrosa DC *et al.* (2015). Endothelial glycocalyx damage is associated with leptospirosis acute kidney injury. *Am J Trop Med Hyg* 92: 611–616.
- Lovell R, Dunn JA, Begum G, Barth NJ, Plant T, Moss PA *et al.* (2005). Soluble syndecan-1 level at diagnosis is an independent prognostic factor in multiple myeloma and the extent of fall from diagnosis to plateau predicts for overall survival. *Br J Haematol* 130: 542–548.
- Luo L, Feng S, Wu Y, Su Y, Jing F, Yi Q (2018a). Serum levels of syndecan-1 in patients with Kawasaki disease. *Pediatr Infect Dis J* : 1.
- Luo Q, Ning P, Zheng Y, Shang Y, Zhou B, Gao Z (2018b). Serum suPAR and syndecan-4 levels predict severity of community-acquired pneumonia: a prospective, multi-centre study. *Crit Care* 22: 15.
- Maeda T, Alexander CM, Friedl A (2004). Induction of syndecan-1 expression in stromal fibroblasts promotes proliferation of human breast cancer cells. *Cancer Res* 64: 612–621.
- Mahtouk K, Hose D, Raynaud P, Hundemer M, Jourdan M, Jourdan E *et al.* (2007). Heparanase influences expression and shedding of syndecan-1, and its expression by the bone marrow environment is a bad prognostic factor in multiple myeloma. *Blood* 109: 4914–4923.
- Maisnar V, Touskova M, Tichy M, Krejsk J, Chrobak L, Voglova J *et al.* (2006). The significance of soluble CD138 in diagnosis of monoclonal gammopathies. *Neoplasma* 53: 26–29.
- Malek-Hosseini Z, Jelodar S, Talei A, Ghaderi A, Doroudchi M (2017). Elevated syndecan-1 levels in the sera of patients with breast cancer correlate with tumor size. *Breast Cancer* 24: 742–747.
- Manon-Jensen T, Itoh Y, Couchman JR (2010). Proteoglycans in health and disease: the multiple roles of syndecan shedding. *FEBS J* 277: 3876–3889.
- Metwaly HA, Al-Gayyar MM, Eletreby S, Ebrahim MA, El-Shishtawy MM (2012). Relevance of serum levels of interleukin-6 and syndecan-1 in patients with hepatocellular carcinoma. *Sci Pharm* 80: 179–188.
- Minowa K, Amano H, Nakano S, Ando S, Watanabe T, Nakiri Y *et al.* (2011). Elevated serum level of circulating syndecan-1 (CD138) in active systemic lupus erythematosus. *Autoimmunity* 44: 357–362.
- Mitselou A, Galani V, Skoufi U, Arvanitis DL, Lampri E, Ioachim E (2016). Syndecan-1, epithelial-mesenchymal transition markers (E-cadherin/ β -catenin) and neoangiogenesis-related proteins (PCAM-1 and endoglin) in colorectal cancer. *Anticancer Res* 36: 2271–2280.
- Mortier A, Van Damme J, Proost P (2012). Overview of the mechanisms regulating chemokine activity and availability. *Immunol Lett* 145: 2–9.
- Mosaad NA, Lotfy HM, Farag YM, Mahfouz RH, Shahin RM (2017). Study of serum syndecan-1 levels in a group of Egyptian juvenile systemic lupus erythematosus patients. *Immunol Lett* 181: 16–19.
- Mukhopadhyay A, Wong MY, Chan SY, Do DV, Khoo A, Ong CT *et al.* (2010). Syndecan-2 and decorin: proteoglycans with a difference--implications in keloid pathogenesis. *J Trauma* 68: 999–1008.
- Mundt F, Heidari-Hamedani G, Nilsonne G, Metintas M, Hjerpe A, Dobra K (2014). Diagnostic and prognostic value of soluble syndecan-1 in pleural malignancies. *Biomed Res Int* 2014: 419853–11.
- Nakao M, Sugaya M, Takahashi N, Otobe S, Nakajima R, Oka T *et al.* (2016). Increased syndecan-4 expression in sera and skin of patients with atopic dermatitis. *Arch Dermatol Res* 308: 655–660.
- Nam EJ, Park PW (2012). Shedding of cell membrane-bound proteoglycans. *Methods Mol Biol* 836: 291–305.
- Nault JC, Guyot E, Laguillier C, Chevret S, Ganne-Carrie N, N'Kontchou G *et al.* (2013). Serum proteoglycans as prognostic biomarkers of hepatocellular carcinoma in patients with alcoholic cirrhosis. *Cancer Epidemiol Biomarkers Prev* 22: 1343–1352.
- Nelson A, Berkestedt I, Schmidtchen A, Ljunggren L, Bodelsson M (2008). Increased levels of glycosaminoglycans during septic shock: relation to mortality and the antibacterial actions of plasma. *Shock* 30: 623–627.
- Neves FM, Meneses GC, Sousa NE, Menezes RR, Parahyba MC, Martins AM *et al.* (2015). Syndecan-1 in acute decompensated heart failure—association with renal function and mortality. *Circ J* 79: 1511–1519.
- Nguyen TH, Liu S, Ong GJ, Stafford I, Frenneaux MP, Horowitz JD (2017). Glycocalyx shedding is markedly increased during the acute phase of Takotsubo cardiomyopathy. *Int J Cardiol* 243: 296–299.
- Nikaido T, Tanino Y, Wang X, Sato S, Misa K, Fukuhara N *et al.* (2015). Serum syndecan-4 as a possible biomarker in patients with acute pneumonia. *J Infect Dis* 212: 1500–1508.
- Ostrowski SR, Haase N, Muller RB, Moller MH, Pott FC, Perner A *et al.* (2015). Association between biomarkers of endothelial injury and hypocoagulability in patients with severe sepsis: a prospective study. *Crit Care* 19: 191.
- Otsuki S, Hanson SR, Miyaki S, Grogan SP, Kinoshita M, Asahara H *et al.* (2010). Extracellular sulfatases support cartilage homeostasis by regulating BMP and FGF signaling pathways. *Proc Natl Acad Sci U S A* 107: 10202–10207.
- Otsuki S, Murakami T, Okamoto Y, Hoshiyama Y, Oda S, Neo M (2017). Suppression of cartilage degeneration by intra-articular injection of heparan sulfate 6-O endosulfatase in a mouse osteoarthritis model. *Histol Histopathol* 32: 725–733.
- Otsuki S, Taniguchi N, Grogan SP, D'Lima D, Kinoshita M, Lotz M (2008). Expression of novel extracellular sulfatases Sulf-1 and Sulf-2 in normal and osteoarthritic articular cartilage. *Arthritis Res Ther* 10: R61.
- Padberg JS, Wiesinger A, di Marco GS, Reuter S, Grabner A, Kentrup D *et al.* (2014). Damage of the endothelial glycocalyx in chronic kidney disease. *Atherosclerosis* 234: 335–343.
- Pap T, Bertrand J (2013). Syndecans in cartilage breakdown and synovial inflammation. *Nat Rev Rheumatol* 9: 43–55.
- Pasqualon T, Pruessmeyer J, Jankowski V, Babendreyer A, Groth E, Schumacher J *et al.* (2015). A cytoplasmic C-terminal fragment of Syndecan-1 is generated by sequential proteolysis and antagonizes Syndecan-1 dependent lung tumor cell migration. *Oncotarget* 6: 31295–31312.
- Pikas DS, Li JP, Vlodavsky I, Lindahl U (1998). Substrate specificity of heparanases from human hepatoma and platelets. *J Biol Chem* 273: 18770–18777.
- Pruessmeyer J, Martin C, Hess FM, Schwarz N, Schmidt S, Kogel T *et al.* (2010). A disintegrin and metalloproteinase 17 (ADAM17) mediates inflammation-induced shedding of syndecan-1 and -4 by lung epithelial cells. *J Biol Chem* 285: 555–564.
- Pumphrey CY, Theus AM, Li S, Parrish RS, Sanderson RD (2002). Neoglycans, carbodiimide-modified glycosaminoglycans: a new class of anticancer agents that inhibit cancer cell proliferation and induce apoptosis. *Cancer Res* 62: 3722–3728.

- Purushothaman A, Chen L, Yang Y, Sanderson RD (2008). Heparanase stimulation of protease expression implicates it as a master regulator of the aggressive tumor phenotype in myeloma. *J Biol Chem* 283: 32628–32636.
- Rapraeger A, Jalkanen M, Bernfield M (1986). Cell surface proteoglycan associates with the cytoskeleton at the basolateral cell surface of mouse mammary epithelial cells. *J Cell Biol* 103: 2683–2696.
- Rapraeger AC, Ell BJ, Roy M, Li X, Morrison OR, Thomas GM *et al.* (2013). Vascular endothelial-cadherin stimulates syndecan-1-coupled insulin-like growth factor-1 receptor and cross-talk between α V β 3 integrin and vascular endothelial growth factor receptor 2 at the onset of endothelial cell dissemination during angiogenesis. *FEBS J* 280: 2194–2206.
- Regos E, Abdelfattah HH, Reszegi A, Szilak L, Werling K, Szabo G *et al.* (2018). Syndecan-1 inhibits early stages of liver fibrogenesis by interfering with TGF β 1 action and upregulating MMP14. *Matrix Biol* 68–69: 474–489.
- Rehm M, Bruegger D, Christ F, Conzen P, Thiel M, Jacob M *et al.* (2007). Shedding of the endothelial glycocalyx in patients undergoing major vascular surgery with global and regional ischemia. *Circulation* 116: 1896–1906.
- Rhiner C, Hengartner MO (2006). Sugar antennae for guidance signals: syndecans and glypicans integrate directional cues for navigating neurons. *ScientificWorldJournal* 6: 1024–1036.
- Rodriguez-Manzanique JC, Carpizo D, Plaza-Calonge Mdel C, Torres-Collado AX, Thai SN, Simons M *et al.* (2009). Cleavage of syndecan-4 by ADAMTS1 provokes defects in adhesion. *Int J Biochem Cell Biol* 41: 800–810.
- Sanaee MN, Malekzadeh M, Khezri A, Ghaderi A, Doroudchi M (2015). Soluble CD138/syndecan-1 increases in the sera of patients with moderately differentiated bladder cancer. *Urol Int* 94: 472–478.
- Sato Y, Tanino Y, Wang X, Nikaido T, Sato S, Misa K *et al.* (2017). Baseline serum syndecan-4 predicts prognosis after the onset of acute exacerbation of idiopathic interstitial pneumonia. *PLoS One* 12: e0176789.
- Saunders S, Jalkanen M, O'Farrell S, Bernfield M (1989). Molecular cloning of syndecan, an integral membrane proteoglycan. *J Cell Biol* 108: 1547–1556.
- Schlondorff J, Blobel CP (1999). Metalloprotease-disintegrins: modular proteins capable of promoting cell-cell interactions and triggering signals by protein-ectodomain shedding. *J Cell Sci* 112 (Pt 21): 3603–3617.
- Schmidt A, Echtermeyer F, Alozie A, Brands K, Buddecke E (2005). Plasmin- and thrombin-accelerated shedding of syndecan-4 ectodomain generates cleavage sites at Lys (114)-Arg (115) and Lys (129)-Val (130) bonds. *J Biol Chem* 280: 34441–34446.
- Schulz JG, Annaert W, Vandekerckhove J, Zimmermann P, De Strooper B, David G (2003). Syndecan 3 intramembrane proteolysis is presenilin/gamma-secretase-dependent and modulates cytosolic signaling. *J Biol Chem* 278: 48651–48657.
- Scudla V, Pika T, Budikova M, Petrova P, Bacovsky J, Srovnalík K *et al.* (2009). The relationship between some soluble osteogenic markers, angiogenic cytokines/other biological parameters and the stages of multiple myeloma evaluated according to the Durie-Salmon and International Prognostic Index stratification systems. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub* 153: 275–282.
- Seidel C, Ringden O, Remberger M (2003). Increased levels of syndecan-1 in serum during acute graft-versus-host disease. *Transplantation* 76: 423–426.
- Seidel C, Sundan A, Hjorth M, Turesson I, Dahl IM, Abildgaard N *et al.* (2000). Serum syndecan-1: a new independent prognostic marker in multiple myeloma. *Blood* 95: 388–392.
- Shah MM, Sakurai H, Gallegos TF, Sweeney DE, Bush KT, Esko JD *et al.* (2011). Growth factor-dependent branching of the ureteric bud is modulated by selective 6-O sulfation of heparan sulfate. *Dev Biol* 356: 19–27.
- Steppan J, Hofer S, Funke B, Brenner T, Henrich M, Martin E *et al.* (2011). Sepsis and major abdominal surgery lead to flaking of the endothelial glycocalyx. *J Surg Res* 165: 136–141.
- Stewart MD, Ramani VC, Sanderson RD (2015). Shed syndecan-1 translocates to the nucleus of cells delivering growth factors and inhibiting histone acetylation: a novel mechanism of tumor-host cross-talk. *J Biol Chem* 290: 941–949.
- Su G, Blaine SA, Qiao D, Friedl A (2007). Shedding of syndecan-1 by stromal fibroblasts stimulates human breast cancer cell proliferation via FGF2 activation. *J Biol Chem* 282: 14906–14915.
- Subramanian SV, Fitzgerald ML, Bernfield M (1997). Regulated shedding of syndecan-1 and -4 ectodomains by thrombin and growth factor receptor activation. *J Biol Chem* 272: 14713–14720.
- Sutherland AE, Sanderson RD, Mayes M, Seibert M, Calarco PG, Bernfield M *et al.* (1991). Expression of syndecan, a putative low affinity fibroblast growth factor receptor, in the early mouse embryo. *Development* 113: 339–351.
- Svennevig K, Kolset SO, Bangstad HJ (2006). Increased syndecan-1 in serum is related to early nephropathy in type 1 diabetes mellitus patients. *Diabetologia* 49: 2214–2216.
- Szarvas T, Reis H, Vom Dorp F, Tschirdewahn S, Niedworok C, Nyirady P *et al.* (2016). Soluble syndecan-1 (SDC1) serum level as an independent pre-operative predictor of cancer-specific survival in prostate cancer. *Prostate* 76: 977–985.
- Takahashi R, Negishi K, Watanabe A, Arai M, Naganuma F, Ohyama Y *et al.* (2011). Serum syndecan-4 is a novel biomarker for patients with chronic heart failure. *J Cardiol* 57: 325–332.
- Torres Filho IP, Torres LN, Salgado C, Dubick MA (2016). Plasma syndecan-1 and heparan sulfate correlate with microvascular glycocalyx degradation in hemorrhaged rats after different resuscitation fluids. *Am J Physiol Heart Circ Physiol* 310: H1468–H1478.
- Tromp J, van der Pol A, Klip IT, de Boer RA, Jaarsma T, van Gilst WH *et al.* (2014). Fibrosis marker syndecan-1 and outcome in patients with heart failure with reduced and preserved ejection fraction. *Circ Heart Fail* 7: 457–462.
- Vassilakopoulos TP, Kyrtsonis MC, Papadogiannis A, Nadali G, Angelopoulou MK, Tzenou T *et al.* (2005). Serum levels of soluble syndecan-1 in Hodgkin's lymphoma. *Anticancer Res* 25: 4743–4746.
- Wang JB, Tian CW, Guo CM, Du HJ, Liu HL, Zhang YJ *et al.* (2008). Increased levels of soluble syndecan-1 in the subretinal fluid and the vitreous of eyes with rhegmatogenous retinal detachment. *Curr Eye Res* 33: 101–107.
- Wang S, Ai X, Freeman SD, Pownall ME, Lu Q, Kessler DS *et al.* (2004). QSulf1, a heparan sulfate 6-O-endosulfatase, inhibits fibroblast growth factor signaling in mesoderm induction and angiogenesis. *Proc Natl Acad Sci U S A* 101: 4833–4838.
- Wang X, Wang H, Yang H, Li J, Cai Q, Shapiro IM *et al.* (2014). Tumor necrosis factor- α - and interleukin-1 β -dependent matrix metalloproteinase-3 expression in nucleus pulposus cells requires cooperative signaling via syndecan 4 and mitogen-activated protein

kinase-NF- κ B axis: implications in inflammatory disc disease. *Am J Pathol* 184: 2560–2572.

Wang XF, Li AM, Li J, Lin SY, Chen CD, Zhou YL *et al.* (2013). Low molecular weight heparin relieves experimental colitis in mice by downregulating IL-1 β and inhibiting syndecan-1 shedding in the intestinal mucosa. *PLoS One* 8: e66397.

Wang Z, Gotte M, Bernfield M, Reizes O (2005). Constitutive and accelerated shedding of murine syndecan-1 is mediated by cleavage of its core protein at a specific juxtamembrane site. *Biochemistry* 44: 12355–12361.

Wu CY, Asano Y, Taniguchi T, Sato S, Yu HS (2016). Serum level of circulating syndecan-1: a possible association with proliferative vasculopathy in systemic sclerosis. *J Dermatol* 43: 63–66.

Yablecovitch D, Oren A, Ben-Horin S, Fudim E, Eliakim R, Saker T *et al.* (2017). soluble syndecan-1: a novel biomarker of small bowel

mucosal damage in children with celiac disease. *Dig Dis Sci* 62: 755–760.

Yang Y, Yaccoby S, Liu W, Langford JK, Pumphrey CY, Theus A *et al.* (2002). Soluble syndecan-1 promotes growth of myeloma tumors in vivo. *Blood* 100: 610–617.

Zhang S, Qing Q, Wang Q, Xu J, Zhi F, Park PW *et al.* (2013). Syndecan-1 and heparanase: potential markers for activity evaluation and differential diagnosis of Crohn's disease. *Inflamm Bowel Dis* 19: 1025–1033.

Zhang Y, Pasparakis M, Kollias G, Simons M (1999). Myocyte-dependent regulation of endothelial cell syndecan-4 expression. Role of TNF- α . *J Biol Chem* 274: 14786–14790.

Zvibel I, Halfon P, Fishman S, Penaranda G, Leshno M, Or AB *et al.* (2009). Syndecan 1 (CD138) serum levels: a novel biomarker in predicting liver fibrosis stage in patients with hepatitis C. *Liver Int* 29: 208–212.